



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	<input checked="" type="radio"/> MIM	Books										
Search	PubMed	<input type="text"/> for					<input type="button" value="Go"/>	<input type="button" value="Clear"/>										
		Limits	Preview/Index	History	Clipboard	Details												
<table border="1"> <tr> <td>Display</td> <td>Abstract</td> <td><input type="button" value="▼"/></td> <td>Show: 20</td> <td><input type="button" value="▼"/></td> <td>Sort</td> <td><input type="button" value="▼"/></td> <td>Send to</td> <td>Text</td> <td><input type="button" value="▼"/></td> </tr> </table>									Display	Abstract	<input type="button" value="▼"/>	Show: 20	<input type="button" value="▼"/>	Sort	<input type="button" value="▼"/>	Send to	Text	<input type="button" value="▼"/>
Display	Abstract	<input type="button" value="▼"/>	Show: 20	<input type="button" value="▼"/>	Sort	<input type="button" value="▼"/>	Send to	Text	<input type="button" value="▼"/>									

Entrez
PubMed

PubMed
Services

Related
Resources

□1: Vaccine 1997 Jun;15(8):853-6

Related Articles, Links

**ELSEVIER SCIENCE
FULL-TEXT ARTICLE**

DNA vaccination for the induction of immune responses against hepatitis C virus proteins.

Inchauspe G, Major ME, Nakano I, Vitvitski L, Trepo C.

INSERM U271, Lyon, France.

Recent analysis of clinical and experimental cases of hepatitis C virus (HCV) infection suggest the possible role of the viral nucleocapsid (C), the nonstructural protein 3 (NS3) and the envelope glycoproteins E1 and/or E2 in the mounting of immune responses capable to control infection (Botarelli et al., Gastroenterology, 1993, 104, 580-587; Choo et al., Proc. Natl Acad. Sci. USA, 1994, 91, 1294-1298). We have used DNA-based immunization to study the immune responses that can be induced by injecting DNA-derived immunogens encoding C and E2 sequences. Comparative analysis were performed in mice using expression plasmids containing full-length or partial gene sequences cloned in fusion with the hepatitis B virus surface antigen (HBV-HCV chimeras). The results obtained indicate that: (1) anti-C and anti-E2 antibodies can be induced with all constructs including the HBV-HCV chimeras; (2) titers range from 1:100 to 1:100000 depending on the antigen and nucleotide sequence context; (3) all HCV DNA immunogens are associated with a predominant Th1 response; (4) CTL can be detected against both HCV and HBV determinants.

PMID: 9234532 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="button" value="▼"/>	Show: 20	<input type="button" value="▼"/>	Sort	<input type="button" value="▼"/>	Send to	Text	<input type="button" value="▼"/>
---------	----------	----------------------------------	----------	----------------------------------	------	----------------------------------	---------	------	----------------------------------

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)